

LINEAR σ -ADDITIVITY AND SOME APPLICATIONS

TAL ORENSHTEIN AND BOAZ TSABAN

ABSTRACT. We show that countable increasing unions preserve a large family of well-studied covering properties, which are not necessarily σ -additive. Using this, together with infinite-combinatorial methods and simple forcing theoretic methods, we explain several phenomena, settle problems of Just, Miller, Scheepers and Szeptycki [15], Gruenhage and Szeptycki [13], Tsaban and Zdomskyy [33], and Tsaban [29, 32], and construct topological groups with very strong combinatorial properties.

1. INTRODUCTION

The following natural definition unifies all results presented here.

Definition 1.1. Let \mathcal{F} be a family of topological spaces. \mathcal{F} is *linearly σ -additive* if it is preserved by countable increasing unions. That is: For each topological space $X = \bigcup_n X_n$ with $X_1 \subseteq X_2 \subseteq \dots$ and $X_n \in \mathcal{F}$ for all $n \in \mathbb{N}$, $X \in \mathcal{F}$.

Removing the restriction that $X_n \subseteq X_{n+1}$ for all n , we obtain the definition of *σ -additive* family. We identify a topological property with the family of all topological spaces satisfying it. Thus, we may talk about linearly σ -additive *properties*.

We consider additivity in the context of topological selection principles, to which we now give a brief introduction.¹ This is a framework suggested by Scheepers in [22] to study in a uniform manner a variety of properties introduced in different mathematical disciplines, since the early 1920's, by Menger, Hurewicz, Rothberger, and Gerlits and Nagy, and many others.

Let X be a topological space. We say that \mathcal{U} is a *cover* of X if $X = \bigcup \mathcal{U}$, but $X \notin \mathcal{U}$. Often, X is considered as a subspace of another space Y , and in this case we always consider covers of X by subsets of Y , and require instead that no member of the cover contains X . Let $\mathcal{O}(X)$ be the family of all open covers of X . Define the following subfamilies of $\mathcal{O}(X)$: $\mathcal{U} \in \Omega(X)$ if each finite subset of X is contained

¹Extended introductions to this field are available in [17, 25, 28].

in some member of \mathcal{U} . $\mathcal{U} \in \Gamma(X)$ if \mathcal{U} is infinite, and each element of X is contained in all but finitely many members of \mathcal{U} .

Some of the following statements may hold for families \mathcal{A} and \mathcal{B} of covers of X .

$S_1(\mathcal{A}, \mathcal{B})$: For all $\mathcal{U}_1, \mathcal{U}_2, \dots \in \mathcal{A}$, there are $U_1 \in \mathcal{U}_1, U_2 \in \mathcal{U}_2, \dots$ such that $\{U_n : n \in \mathbb{N}\} \in \mathcal{B}$.

$S_{\text{fin}}(\mathcal{A}, \mathcal{B})$: For all $\mathcal{U}_1, \mathcal{U}_2, \dots \in \mathcal{A}$, there are finite $\mathcal{F}_1 \subseteq \mathcal{U}_1, \mathcal{F}_2 \subseteq \mathcal{U}_2, \dots$ such that $\bigcup_n \mathcal{F}_n \in \mathcal{B}$.

$U_{\text{fin}}(\mathcal{A}, \mathcal{B})$: For all $\mathcal{U}_1, \mathcal{U}_2, \dots \in \mathcal{A}$, none containing a finite subcover, there are finite $\mathcal{F}_1 \subseteq \mathcal{U}_1, \mathcal{F}_2 \subseteq \mathcal{U}_2, \dots$ such that $\{\bigcup \mathcal{F}_n : n \in \mathbb{N}\} \in \mathcal{B}$.

We say, e.g., that X satisfies $S_1(O, O)$ if the statement $S_1(O(X), O(X))$ holds. This way, $S_1(O, O)$ is a property of topological spaces, and similarly for all other statements and families of covers. Under some mild hypotheses on the considered topological spaces, each nontrivial property among these properties, where \mathcal{A}, \mathcal{B} range over O, Ω, Γ , is equivalent to one in Figure 1 [22, 15]. In this diagram, an arrow denotes implication.




FIGURE 1. The Scheepers Diagram

In this diagram, the classical name of a property is indicated below it, as well as two names ending with a symbol \uparrow , by which we indicate that the properties $S_1(\Omega, \Omega)$ and $S_{\text{fin}}(\Omega, \Omega)$ may also be viewed as classical ones [21, 15], in accordance with the following notation.

Definition 1.2. Let P be a property of topological spaces. X satisfies P^\uparrow if all finite powers X^k of X satisfy P .

The Scheepers diagram is at the heart of the field of topological selection principles, and many additional—classical and new—properties are studied in relation to it. The reader is encouraged to consider his favorite properties in light of the results presented here.

In Section 2, we prove that all properties in the Scheepers Diagram are linearly σ -additive, and are thus hereditary for F_σ subsets. This solves a problem of Tsaban and Zdomskyy from [33].

A crucial part of the proof that the studied properties are linearly σ -additive, is a recent theorem of F. Jordan [14]. Miller asked in [20] whether $S_1(\Omega, \Gamma)$ is linearly σ -additive. A negative solution would have solved the notorious Gerlits-Nagy problem [12]. Using a brilliant argument, Jordan proved that this is not the case. We give a direct version of Jordan's solution, and provide some applications.

In Section 3, we use Jordan's method in a proof that if there is an unbounded family of cardinality \aleph_1 in the Baire space, then there is an uncountable set of reals satisfying $S_1(\Omega, \Gamma)$. This settles in the positive of a problem from the seminal paper of Just, Miller, Scheepers, and Szeptycki [15]. Indeed, our result is more general, and also solves a problem of Gruenhage and Szeptycki [13].

In Section 4, we apply linear σ -additivity to study heredity of properties, answer a question of Zdomskyy, and suggest a simple revision of a question of Bukovský, Reclaw, and Repický, which makes it possible to answer it in the positive. (The problem, as originally stated, was answered in the negative by Miller [18].) This section also explains a phenomenon observed in [2].

In Section 5 we apply our results to construct topological groups with strong combinatorial properties, and solve problems from [29] and [32].

1.1. Generalizations. The results presented here require little or no assumptions on the topology of the studied spaces. However, they are interesting even when restricting attention to, e.g., metric spaces or even subsets of \mathbb{R} .

For concreteness, we present the results only for the types of covers mentioned above, but the proofs show that they hold for many additional types. In particular, define B , B_Ω , B_Γ as O , Ω , Γ were defined, replacing *open cover* by *countable Borel cover*. The properties thus obtained have rich history of their own [26]. Mild assumptions on X imply that the considered open covers may be assumed to be countable, and this makes the Borel variants of the studied properties (strictly) stronger [26]. All of the results presented here also hold in the Borel case (after replacing *open* or *closed* by *Borel*). Moreover, unlike some of the results in the open case, none of the Borel variant requires any assumption on the topology of X .

2. LINEAR σ -ADDITIVITY IN THE SCHEEPERS DIAGRAM

One motivation for studying linear σ -additivity in the context of the Scheepers Diagram is an experimentally observed dichotomy concerning additivity in of properties in this diagram: Each property there is either provably σ -additive, or else not even provably *finitely* additive.² In this section, we prove the following.

Theorem 2.1. *All properties in the Scheepers Diagram (Figure 1) are linearly σ -additive.*

It remains to notice the following.

Lemma 2.2. *Each linearly σ -additive property is either σ -additive, or else not finitely additive.*

Proof. $\bigcup_n X_n = \bigcup_n (\bigcup_{m \leq n} X_m)$. \square

Corollary 2.3. *Each property in the Scheepers Diagram is either σ -additive, or else not additive.* \square

Before proving Theorem 2.1, we point out two additional consequences. In Problem 4.9 of [33] and Problem 6.2 of [32], Tsaban and Zdomskyy ask whether $S_1(\Gamma, \Omega)$ and $S_{\text{fin}}(\Gamma, \Omega)$ are hereditary for F_σ subsets. We obtain a positive answer.

Corollary 2.4. *All properties in the Scheepers Diagram are hereditary for F_σ subsets.*

Proof. These properties are easily seen to be hereditary for closed subsets [15], and countable unions of closed subsets can be presented as countable increasing unions of closed subsets. Apply Theorem 2.1. \square

The following was observed in the past for at least some properties in the Scheepers Diagram, each time using an ad-hoc argument.

Corollary 2.5. *Let P be a property in the Scheepers Diagram, and X, D be subspaces of some topological space. If X satisfies P and D is countable, then $X \cup D$ satisfies P .*

Proof. In light of Theorem 2.1, it remains to observe that for each singleton $\{a\}$, $X \cup \{a\}$ satisfies P . This is not hard to verify. \square

Theorem 2.1 is proved in parts. The properties $S_1(O, O)$, $S_1(\Gamma, O)$, $S_{\text{fin}}(O, O)$, $S_1(\Gamma, \Gamma)$, and $U_{\text{fin}}(O, \Gamma)$ are in fact σ -additive [15, 24] (see [30]).

Linear σ -additivity of $U_{\text{fin}}(O, \Omega)$ was proved in [33] for sets of reals. It also follows from the following.

²A survey of the involved results, with complete proofs, is available in [30].

Theorem 2.6. *For all $\Pi \in \{S_1, S_{\text{fin}}, U_{\text{fin}}\}$ and $\mathcal{A} \in \{\Gamma, \Omega, O\}$, $\Pi(\mathcal{A}, \Omega)$ is linearly σ -additive.*

Proof. We prove the theorem for $\Pi = S_{\text{fin}}$; the other proofs being similar.

Assume that $X = \bigcup_n X_n$ is an increasing union, with each X_n satisfying $S_{\text{fin}}(\mathcal{A}, \Omega)$. Let $\mathcal{U}_1, \mathcal{U}_2, \dots \in \mathcal{A}(X)$. We first exclude the trivial case: Assume that for infinitely many n , there are m_n and elements $U_{m_n} \in \mathcal{U}_{m_n}$ such that $X_n \subseteq U_{m_n}$. As X is not contained in any member of any \mathcal{U}_n and the sets X_n increase to X , we may if necessary thin out the sequence m_n to make it increasing. Then $\{U_{m_n} : n \in \mathbb{N}\} \in \Omega(X)$, and this suffices.

Thus, we may assume that for all n, k , we have that $\mathcal{U}_k \in \mathcal{A}(X_n)$. Take a partition $\mathbb{N} = \bigcup_n I_n$ of \mathbb{N} into infinite sets I_n . Fix n . As X_n satisfies $S_{\text{fin}}(\mathcal{A}, \Omega)$, there are finite $\mathcal{F}_k \subseteq \mathcal{U}_k$, $k \in I_n$, such that $\bigcup_{k \in I_n} \mathcal{F}_k \in \Omega(X_n)$. Then $\bigcup_{k \in \mathbb{N}} \mathcal{F}_k \in \Omega(X)$. \square

The remaining property, $S_1(\Omega, \Gamma)$, was treated by F. Jordan.

2.1. Jordan's Theorem and some applications. The following technical lemma will be useful in the proof of Jordan's Theorem below.

Lemma 2.7. *Let $Y \subseteq X$ be such that Y satisfies $S_1(\Gamma, \Gamma)$. Assume that for each n ,*

- (1) \mathcal{U}_n is an infinite family of open subsets of X ; and
- (2) For each $y \in Y$, $y \in U$ for all but finitely many $U \in \mathcal{U}_n$.

Then there are infinite $\mathcal{V}_1 \subseteq \mathcal{U}_1, \mathcal{V}_2 \subseteq \mathcal{U}_2, \dots$, such that for each $y \in Y$, $y \in \bigcap \mathcal{V}_n$ for all but finitely many n .

Proof. It may be the case that no subset of \mathcal{U}_n is in $\Gamma(Y)$.

Case 1: For all but finitely many n , $\mathcal{V}_n = \{U \in \mathcal{U}_n : Y \subseteq U\}$ is infinite. Then the sets \mathcal{V}_n thus defined are as required.

Case 2: Let I be the set of all n such that $\mathcal{V}_n = \{U \in \mathcal{U}_n : Y \subseteq U\}$ is infinite, and $J = \mathbb{N} \setminus I$. For each $n \in J$, $\mathcal{W}_n = \{U \in \mathcal{U}_n : Y \not\subseteq U\}$ is infinite, and thus $\mathcal{W}_n \in \Gamma(Y)$. As J is infinite, there are by Theorem 15 of [31] infinite $\mathcal{V}_n \subseteq \mathcal{W}_n$, $n \in J$, such that each $y \in Y$ belongs to $\bigcap \mathcal{V}_n$ for all but finitely many n . (Briefly: By thinning out if needed, we may assume that each \mathcal{W}_n is countable and that $\mathcal{W}_n \cap \mathcal{W}_m = \emptyset$ for $m \neq n$ [22]. Apply $S_1(\Gamma, \Gamma)$ to the countable family of all cofinite subsets of all \mathcal{W}_n to obtain $\mathcal{V} \in \Gamma(Y)$. Let $\mathcal{V}_n = \mathcal{V} \cap \mathcal{W}_n$.) The sets $\mathcal{V}_1, \mathcal{V}_2, \dots$ are as required. \square

Theorem 2.8 (Jordan [14]). $S_1(\Omega, \Gamma)$ is linearly σ -additive.

Proof. We give a direct proof, following what seems to be the essence of Jordan's arguments. The following statement can be deduced from Theorem 7 in [14].

Lemma 2.9. *Let $X = \bigcup_n X_n$ be an increasing union, where each X_n satisfies $S_1(\Gamma, \Gamma)$. For all $\mathcal{U}_1 \in \Gamma(X_1), \mathcal{U}_2 \in \Gamma(X_2), \dots$, there are infinite $\mathcal{V}_1 \subseteq \mathcal{U}_1, \mathcal{V}_2 \subseteq \mathcal{U}_2, \dots$, such that for each $x \in X$, $x \in \bigcap \mathcal{V}_n$ for all but finitely many n .*

Proof. Step 1: By Lemma 2.7, we may thin out the families \mathcal{U}_n , so that they remain infinite, and each member of X_1 belongs to all but finitely many $\bigcap \mathcal{U}_n$. Let $\mathcal{V}_1 = \mathcal{U}_1$.

Step 2: By the same Lemma, we may thin out further the families $\mathcal{U}_n, n \geq 2$, so that they remain infinite, and each member of X_2 belongs to all but finitely many $\bigcap \mathcal{U}_n, n > 1$. Let $\mathcal{V}_2 = \mathcal{U}_2$.

Step k : By Lemma 2.7, we may thin out further the families $\mathcal{U}_n, n \geq k$, so that they remain infinite, and each member of X_k belongs to all but finitely many $\bigcap \mathcal{U}_n, n \geq k$. Let $\mathcal{V}_k = \mathcal{U}_k$.

The sets $\mathcal{V}_1, \mathcal{V}_2, \dots$ are as required. \square

Now, let $X = \bigcup_n X_n$ be an increasing union, where each X_n satisfies $S_1(\Omega, \Gamma)$. Let $\mathcal{U}_1, \mathcal{U}_2, \dots \in \Omega(X)$. As in the argument of the proof of Lemma 2.7, we may assume that $\mathcal{U}_n \in \Omega(X_n)$ for all n . As X_n satisfies $S_1(\Omega, \Gamma)$, we may thin out \mathcal{U}_n so that $\mathcal{U}_n \in \Gamma(X_n)$.³ Apply Lemma 2.9, and pick for each n some $U_n \in \mathcal{V}_n \setminus \{U_1, \dots, U_{n-1}\}$. $\{U_n : n \in \mathbb{N}\} \in \Gamma(X)$. \square

Let $C(X)$ be the family of continuous functions $f : X \rightarrow \mathbb{R}$. $C(X)$ has the *Arhangel'skii's property* α_2 if the following holds for all $f_m^n \in C(X), n, m \in \mathbb{N}$: If for each $n, \lim_{m \rightarrow \infty} f_m^n(x) = 0$ for all $x \in X$, then there are m_n such that $\lim_n f_{m_n}^n(x) = 0$ for each $x \in X$.

For the same reason briefly mentioned in the proof of Lemma 2.7, one may require in the definition of α_2 that there are infinite $I_1, I_2, \dots \subseteq \mathbb{N}$, such that some (equivalently, any) enumeration of the countable set $\bigcup_n \{f_m^n(x) : m \in I_n\}$ converges to 0, for each $x \in X$. Indeed, this was Arhangel'skii's original definition of α_2 .

Definition 2.10. Let $X = \bigcup_n X_n$ be an increasing union. $C(X)$ is $\{X_n\}_{n \in \mathbb{N}}\text{-}\alpha_2$ if the following holds whenever $f_m^n \in C(X_n)$ for all m, n : If for each $n, \lim_m f_m^n(x) = 0$ for all $x \in X_n$, then there are infinite

³Clearly, if $\mathcal{U} \in \Omega(X)$ and X satisfies $S_1(\Omega, \Gamma)$, then \mathcal{U} contains a subcover \mathcal{V} such that $\mathcal{V} \in \Gamma(X)$ [12].

$I_1, I_2, \dots \subseteq \mathbb{N}$, such that for each $x \in X$, $\lim_n \sup\{|f_m^n(x)| : m \in I_n\} = 0$.⁴

Clearly, $\{X_n\}_{n \in \mathbb{N}}\text{-}\alpha_2$ implies α_2 . The proof of Lemma 2.9, with minor modifications, yields the following new result about function spaces.

Theorem 2.11. *Let $X = \bigcup_n X_n$ be an increasing union, such that $C(X_n)$ is α_2 for all n . Then $C(X)$ is $\{X_n\}_{n \in \mathbb{N}}\text{-}\alpha_2$.*

Proof. Assume that $f_m^n \in C(X_n)$ for all m, n , and for each n , $\lim_m f_m^n(x) = 0$ for all $x \in X_n$.

Step 1: As $C(X_1)$ is α_2 , there are infinite $J_1, J_2, \dots \subseteq \mathbb{N}$ such that any enumeration of the countable set $\bigcup_n \{f_m^n(x) : m \in J_n\}$ converges to 0, for each $x \in X_1$. In particular, $\lim_n \sup\{|f_m^n(x)| : m \in J_n\} = 0$ for all $x \in X_1$. Let $I_1 = J_1$.

Step k : For each $n \geq k$, $\lim_{m \in J_n} f_m^n(x) = 0$ for all $x \in X_k$. As $C(X_k)$ is α_2 , the index sets J_k, J_{k+1}, \dots may be thinned out so that they remain infinite, and $\lim_n \sup\{|f_m^n(x)| : m \in J_n\} = 0$ for all $x \in X_k$. Let $I_k = J_k$.

The index sets I_1, I_2, \dots are as required. \square

Corollary 2.12. *Assume that for each subset Y of X , $C(Y)$ is α_2 . For all $X_1 \subseteq X_2 \subseteq \dots \subseteq X$, $C(\bigcup_n X_n)$ is $\{X_n\}_{n \in \mathbb{N}}\text{-}\alpha_2$.* \square

A topological space is *perfectly normal* if for each pair of disjoint closed sets $C_0, C_1 \subseteq X$, there is $f \in C(X)$ such that $f^{-1}[\{0\}] = C_0$ and $f^{-1}[\{1\}] = C_1$.

Corollary 2.13. *Assume that X is a perfectly normal space and $C(X)$ is α_2 . For all closed sets $X_1 \subseteq X_2 \subseteq \dots \subseteq X$, $C(\bigcup_n X_n)$ is $\{X_n\}_{n \in \mathbb{N}}\text{-}\alpha_2$.*

Proof. If $C(X)$ is α_2 and Y is a closed subset of X , then $C(Y)$ is α_2 .⁵ Apply Theorem 2.11. \square

In addition to these applications, Jordan's Lemma 2.9 is an ingredient in a solution of a thus far open problem, to which we now turn.

⁴For each $x \in X$, $x \in X_n$ for all large enough n . Thus, for all large enough n , $r_n(x) = \sup\{|f_m^n(x)| : m \in I_n\}$ is defined, and therefore the question, whether $\lim_n r_n(x)$ converges to 0 or not, makes sense.

⁵The argument is as in Theorem 4.1 of [6]: as X is perfectly normal, the open set $X \setminus Y$ is a countable increasing union of closed sets, $X \setminus Y = \bigcup_n C_n$. For each n , extend each f_m^n to an element of $C(X)$ which is constantly 0 on C_n . Applying α_2 on the new sequences, we obtain m_n such that $\lim_n f_{m_n}^n(x) = 0$ for all $x \in X$, and thus this is also the case for the original functions, for all $x \in Y$.

3. γ -SETS FROM A WEAK HYPOTHESIS

In this section, we construct sets of reals satisfying $S_1(\Omega, \Gamma)$, traditionally called γ -sets, from a weak set theoretic hypothesis. The problem thus settled has some history, which we now survey briefly. This involves combinatorial cardinal characteristics of the continuum [3]. We give the necessary definitions as we proceed. Readers who are new to this field, may skip this section in their first reading.

γ -sets were introduced by Gerlits and Nagy in [12], their most influential paper, as the third property in a list numbered α through ϵ . This turned out to be the most important property in the list, and obtained its item number as its name. One of the main results in [12] is that for Tychonoff spaces X , $C(X)$ with the topology of pointwise convergence is Fréche-Urysohn if, and only if, X is a γ -set.

While uncountable γ -sets exist in ZFC,⁶ Borel's Conjecture (which is consistent with, but not provable within, ZFC) implies that all metrizable γ -sets are countable.

Since we are dealing with constructions rather than general results, we restrict attention in this section to subsets of \mathbb{R} (or, since the property is preserved by continuous images, subsets of any topological space which can be embedded in \mathbb{R}). Thus, we may restrict attention to *countable* open covers.

Gerlits and Nagy proved in [12] that Maritn's Axiom implies that all spaces of cardinality less than \mathfrak{c} are γ -sets. There is a simple reason for that: The *critical cardinality* of a property P , denoted $\text{non}(P)$, is the minimal cardinality of a set not satisfying P . Let $(\frac{\Omega}{\Gamma})$ be the property: Each $\mathcal{U} \in \Omega(X)$ contains a set $\mathcal{V} \in \Gamma(X)$. Gerlits and Nagy proved that $S_1(\Omega, \Gamma) = (\frac{\Omega}{\Gamma})$ [12]. Let $A \subseteq^* B$ mean that $A \setminus B$ is finite. A is a *pseudointersection* of \mathcal{F} if $A \subseteq^* B$ for all $B \in \mathcal{F}$. Let \mathfrak{p} be the minimal cardinality of a family \mathcal{F} of infinite subsets of \mathbb{N} which is closed under finite intersections, and has no pseudointersection. Then $\text{non}(\frac{\Omega}{\Gamma}) = \mathfrak{p}$ [11], and Maritn's Axiom implies $\mathfrak{p} = \mathfrak{c}$ [11].

By definition, for each property P , every space of cardinality smaller than $\text{non}(P)$ satisfies P . Thus, the real question is whether there is X of cardinality at least $\text{non}(P)$, which satisfies P . Galvin and Miller [11] proved a result of this type: $\mathfrak{p} = \mathfrak{c}$ implies that there is a γ -set of reals, of cardinality \mathfrak{p} . Just, Miller, Scheepers and Szeptycki [15] have improved the construction of [11]. We introduce their construction in a slightly more general form, that will be useful later.

⁶The axioms of Zermelo and Fraenkel, together with the axiom of Choice, the ordinary axioms of mathematics.

Cantor's space $\{0, 1\}^{\mathbb{N}}$ is equipped with the Tychonoff product topology, and $P(\mathbb{N})$ is identified with $\{0, 1\}^{\mathbb{N}}$ using characteristic functions. This defines the topology of $P(\mathbb{N})$. The partition $P(\mathbb{N}) = [\mathbb{N}]^{\infty} \cup [\mathbb{N}]^{<\infty}$, into the infinite and the finite sets, respectively, is useful here.

For $f, g \in \mathbb{N}^{\mathbb{N}}$, let $f \leq^* g$ if $f(n) \leq g(n)$ for all but finitely many n . \mathfrak{b} is the minimal cardinality of a \leq^* -unbounded subset of $\mathbb{N}^{\mathbb{N}}$. A set $B \subseteq [\mathbb{N}]^{\infty}$ is *unbounded* if the set of all increasing enumerations of elements of B is unbounded in $\mathbb{N}^{\mathbb{N}}$, with respect to \leq^* . It follows that $|B| \geq \mathfrak{b}$. For $m, n \in \mathbb{N}$, let $(m, n) = \{k : m < k < n\}$.

Lemma 3.1 (folklore). *If $B \subseteq [\mathbb{N}]^{\infty}$ is unbounded, then for each increasing $f \in \mathbb{N}^{\mathbb{N}}$, there is $x \in B$ such that $x \cap (f(n), f(n+1)) = \emptyset$ for infinitely many n .*

Proof. Assume that f is a counterexample. Let g dominate all functions $f_m(n) = f(n+m)$, $m \in \mathbb{N}$. Then for each $x \in B$, $x \leq^* g$. Indeed, let m be such that for all $n \geq m$, $x \cap (f(n), f(n+1)) \neq \emptyset$. Then for each n , the n -th element of x is smaller than $f_{m+1}(n)$. \square

Definition 3.2. A *tower* of cardinality κ is a set $T \subseteq [\mathbb{N}]^{\infty}$ which can be enumerated bijectively as $\{x_{\alpha} : \alpha < \kappa\}$, such that for all $\alpha < \beta < \kappa$, $x_{\beta} \subseteq^* x_{\alpha}$.

An *unbounded tower* of cardinality κ is an unbounded set $T \subseteq [\mathbb{N}]^{\infty}$ which is a tower of cardinality κ . (Necessarily, $\kappa \geq \mathfrak{b}$.)

Let \mathfrak{t} be the minimal cardinality of a tower which has no pseudointersection. Rothberger proved that $\mathfrak{t} \leq \mathfrak{b}$ [3].

Lemma 3.3 (folklore). *$\mathfrak{t} = \mathfrak{b}$ if, and only if, there is an unbounded tower of cardinality \mathfrak{t} .*

Proof. (\Rightarrow) Construct x_{α} by induction on α . Let $\{b_{\alpha} : \alpha < \mathfrak{b}\} \subseteq \mathbb{N}^{\mathbb{N}}$ be unbounded. At step α , let a be a pseudointersection of $\{x_{\beta} : \beta < \alpha\}$, and take $x_{\alpha} \subseteq a$ such that the increasing enumeration of x_{α} dominates b_{α} .

(\Leftarrow) $\mathfrak{t} \leq \mathfrak{b} \leq |T| = \mathfrak{t}$. \square

Just, Miller, Scheepers and Szeptycki [15] proved that if T is an unbounded tower of cardinality \aleph_1 , then $T \cup [\mathbb{N}]^{<\infty}$ satisfies $\mathbf{S}_1(\Omega, \Omega)$, as well as a property, which was later proved by Scheepers [24] to be equivalent to $\mathbf{S}_1(\Gamma, \Gamma)$. In Problem 7 of [15], we are asked the following.

Problem 3.4 (Just-Miller-Scheepers-Szeptycki [15]). *Assume that $T \subseteq [\mathbb{N}]^{\infty}$ is an unbounded tower of cardinality \aleph_1 (so that $\aleph_1 = \mathfrak{b}$). Is $T \cup [\mathbb{N}]^{<\infty}$ a γ -set, i.e., satisfies $\mathbf{S}_1(\Omega, \Gamma)$?*

Scheepers proves in [23] that for each unbounded tower T of cardinality $\mathfrak{t} = \mathfrak{b}$, $T \subseteq [\mathbb{N}]^\infty$ satisfies $S_1(\Gamma, \Gamma)$.

Miller [19] proves that in the Hechler model, there are no uncountable γ -sets. In this model, $\aleph_1 = \mathfrak{p} = \mathfrak{t} < \mathfrak{b}$, and thus $\aleph_1 = \mathfrak{t}$ does not suffice to have an uncountable γ -set of reals. At the end of [19] and in its appendix, Miller proves that $\Diamond(\mathfrak{b})$, a property strictly stronger than $\aleph_1 = \mathfrak{b}$, implies that there is an uncountable γ -set of reals.⁷ He concludes that it is still open whether $\mathfrak{b} = \aleph_1$ is enough to construct an uncountable γ -set.

We show that the answer is positive, and indeed also answer a question of Gruenhage and Szeptycki [13]: A classical problem of Malykhin asks whether there is a countable Fréchet-Urysohn topological group which is not metrizable. Gruenhage and Szeptycki prove that $F \subseteq \mathbb{N}^\mathbb{N}$ is a γ -set if, and only if, a certain construction associated to F provides a positive answer to Malykhin's Problem [13]. They define a generalization of γ -set, called *weak γ -set*, and combine their results with results of Nyikos to prove that $\mathfrak{p} = \mathfrak{b}$ implies that there is a weak γ -set in $\mathbb{N}^\mathbb{N}$ [13, Corollary 10]. They write: "The relationship between γ -sets and weak γ -sets is not known. Perhaps $\mathfrak{b} = \mathfrak{p}$ implies the existence of a γ -set." Our solution confirms their conjecture.

$\mathfrak{p} \leq \mathfrak{t} \leq \mathfrak{b}$, and in all known models of set theory, $\mathfrak{p} = \mathfrak{t}$. Our theorem reproduces Galvin and Miller's Theorem when $\mathfrak{p} = \mathfrak{c}$ [11], but gives additional information: Even if the possible open covers are not considered during the construction in [11], the resulting set is still a γ -set.

Theorem 3.5. *For each unbounded tower T of cardinality \mathfrak{p} in $[\mathbb{N}]^\infty$, $T \cup [\mathbb{N}]^{<\infty}$ satisfies $S_1(\Omega, \Gamma)$.*

Proof. By Lemma 3.3, there is an unbounded tower T of cardinality \mathfrak{p} if, and only if, $\mathfrak{p} = \mathfrak{b}$. Let $T = \{x_\alpha : \alpha < \mathfrak{b}\}$ be an unbounded tower of cardinality \mathfrak{b} . For each α , let $X_\alpha = \{x_\beta : \beta < \alpha\} \cup [\mathbb{N}]^{<\infty}$. We will show that $T \cup [\mathbb{N}]^{<\infty}$ satisfies $\left(\frac{\Omega}{\Gamma}\right)$. Let $\mathcal{U} \in \Omega(T \cup [\mathbb{N}]^{<\infty})$. We use the following modification of Lemma 1.2 of [11].

Lemma 3.6. *Assume that $[\mathbb{N}]^{<\infty} \subseteq X \subseteq P(\mathbb{N})$ and X satisfies $\left(\frac{\Omega}{\Gamma}\right)$. For each $\mathcal{U} \in \Omega(X)$, there are $m_1 < m_2 < \dots$ and distinct $U_1, U_2, \dots \in \mathcal{U}$ such that $\{U_n : n \in \mathbb{N}\} \in \Gamma(X)$, and for each $x \subseteq \mathbb{N}$, $x \in U_n$ whenever $x \cap (m_n, m_{n+1}) = \emptyset$.*

Proof. As X satisfies $\left(\frac{\Omega}{\Gamma}\right)$, we may thin out \mathcal{U} so that $\mathcal{U} \in \Gamma(X)$.

We proceed as in the proof of Lemma 1.2 of [11]. Let $m_1 = 1$. For each $n \geq 1$: As $\mathcal{U} \in \Omega(X)$, each finite subset of X is contained in

⁷ $\Diamond(\mathfrak{b})$ is defined in Dzamonja-Hrusak-Moore [9].

infinitely many elements of \mathcal{U} . Take $U_n \in \mathcal{U} \setminus \{U_1, \dots, U_{n-1}\}$, such that $P(\{1, \dots, m_n\}) \subseteq U_n$. As U_n is open, for each $s \subseteq \{1, \dots, m_n\}$ there is k_s such that for each $x \in P(\mathbb{N})$ with $x \cap \{1, \dots, k_s - 1\} = s$, $x \in U_n$. Let $m_{n+1} = \max\{k_s : s \subseteq \{1, \dots, m_n\}\}$.

As $\{U_n : n \in \mathbb{N}\}$ is an infinite subset of $\mathcal{U} \in \Gamma(X)$, $\{U_n : n \in \mathbb{N}\} \in \Gamma(X)$, too. \square

We may assume that \mathcal{U} is countable. As \mathfrak{b} is regular, there is $\alpha_1 < \mathfrak{b}$ such that X_{α_1} is not contained in any member of \mathcal{U} . This guarantees that $\mathcal{U} \in \Omega(X_\alpha)$ for all $\alpha \geq \alpha_1$.

As $|X_{\alpha_1}| < \mathfrak{p}$, X_{α_1} satisfies $(\frac{\Omega}{\Gamma})$. As $\mathcal{U} \in \Omega(X_{\alpha_1})$, by Lemma 3.6 there are $m_1^1 < m_2^1 < \dots$ and distinct $U_1^1, U_2^1, \dots \in \mathcal{U}$ such that $\{U_n^1 : n \in \mathbb{N}\} \in \Gamma(X_{\alpha_1})$, and for each $x \in P(\mathbb{N})$, $x \in U_n^1$ whenever $x \cap (m_n^1, m_{n+1}^1) = \emptyset$. Let $D_1 = \mathbb{N}$.

As $\alpha_1 < \mathfrak{b}$, $\{x_\alpha : \alpha_1 < \alpha < \mathfrak{b}\}$ is unbounded. By Lemma 3.1, there is $\alpha_2 > \alpha_1$ such that $D_2 = \{n : x_{\alpha_2} \cap (m_n^1, m_{n+1}^1) = \emptyset\}$ is infinite. As $|X_{\alpha_2}| < \mathfrak{p}$, X_{α_2} satisfies $(\frac{\Omega}{\Gamma})$. As $\mathcal{U} \in \Omega(X_{\alpha_2})$, by Lemma 3.6 there are $m_1^2 < m_2^2 < \dots$ and distinct $U_1^2, U_2^2, \dots \in \mathcal{U}$ such that $\{U_n^2 : n \in \mathbb{N}\} \in \Gamma(X_{\alpha_2})$, and for each $x \in P(\mathbb{N})$, $x \in U_n^2$ whenever $x \cap (m_n^2, m_{n+1}^2) = \emptyset$. As D_2 is infinite, $\{U_n^2 : n \in D_2\} \in \Gamma(X_{\alpha_2})$.

Continue in the same manner, to define for each $k > 1$ elements with the following properties:

- (1) $\alpha_k > \alpha_{k-1}$;
- (2) $D_k = \{n : x_{\alpha_k} \cap (m_n^{k-1}, m_{n+1}^{k-1}) = \emptyset\}$ is infinite;
- (3) $m_1^k < m_2^k < \dots$;
- (4) $U_1^k, U_2^k, \dots \in \mathcal{U}$ are distinct;
- (5) $\{U_n^k : n \in D_k\} \in \Gamma(X_{\alpha_k})$; and
- (6) For each $x \in P(\mathbb{N})$, $x \in U_n^k$ whenever $x \cap (m_n^k, m_{n+1}^k) = \emptyset$.

Let $\alpha = \sup_k \alpha_k$. As \mathfrak{b} is regular, $\alpha < \mathfrak{b}$. $X_\alpha = \bigcup_k X_{\alpha_k}$ is a countable increasing union. For each k , $|X_{\alpha_k}| < \mathfrak{b}$, and thus X_{α_k} satisfies $\mathbf{S}_1(\Gamma, \Gamma)$. By Lemma 2.9, there are infinite $I_1 \subseteq D_1, I_2 \subseteq D_2, \dots$ such that each $x \in X_\alpha$ belongs to $\bigcap_{n \in I_k} U_n^k$ for all but finitely many $k \in \mathbb{N}$.

Take $n_1 \in I_2$. For $k > 1$, take $n_k \in I_{k+1}$ such that $m_{n_k}^k > m_{n_{k-1}+1}^{k-1}$, $x_\alpha \cap (m_{n_k}^k, m_{n_k+1}^k) \subseteq x_{\alpha_{k+1}} \cap (m_{n_k}^k, m_{n_k+1}^k)$, and $U_{n_k}^k \notin \{U_{n_1}^1, \dots, U_{n_{k-1}}^{k-1}\}$. We claim that $\{U_{n_k}^k : k \in \mathbb{N}\} \in \Gamma(T \cup [\mathbb{N}]^{<\infty})$. As $\{U_{n_k}^k : k \in \mathbb{N}\} \in \Gamma(X_\alpha)$, it remains to show that for each $x \subseteq^* x_\alpha$, $x \in U_{n_k}^k$ for all but finitely many k . For each large enough k , $m_{n_k}^k$ is large enough, so that

$$x \cap (m_{n_k}^k, m_{n_k+1}^k) \subseteq x_\alpha \cap (m_{n_k}^k, m_{n_k+1}^k) \subseteq x_{\alpha_{k+1}} \cap (m_{n_k}^k, m_{n_k+1}^k) = \emptyset,$$

since $n_k \in D_{k+1}$. Thus, $x \in U_{n_k}^k$. \square

Remark 3.7. Zdomskyy points out that our proof actually shows that a wider family of sets are γ -sets. For example, if we start with T an unbounded tower of cardinality \mathfrak{p} , and thin out its elements arbitrarily, $T \cup [\mathbb{N}]^{<\omega}$ remains a γ -set. This may be useful for constructions of examples with additional properties, since this way, each element of T may be chosen arbitrarily from a certain perfect set.

In particular, we have that in each model of ZFC where $\mathfrak{p} = \mathfrak{b}$, there are γ -sets of cardinality \mathfrak{p} .

Corollary 3.8. *In each of the Cohen, Random, Sacks, and Miller models of ZFC, there are γ -sets of reals with cardinality \mathfrak{p} .* \square

Earlier, Corollary 3.8 was shown for the Sacks model by Ciesielski, Millán, and Pawlikowski in [8], and for the Cohen and Miller models by Miller [19], using specialized arguments. It seems that the result, that there are uncountable γ -sets in the Random reals model (constructed by extending a model of the Continuum Hypothesis), is new.

As discussed above, there are no uncountable γ -sets in the Hechler model [19]. Since the Laver and Mathias models satisfy Borel's Conjecture, there are no uncountable γ -sets in these models, too.

4. HEREDITY

A topological space X satisfies a property P *hereditarily* if each subspace of X satisfies P . In our context, heredity was observed to be tightly connected to the following property. X is a σ -space if each Borel subset of X is an F_σ subset of X .

A combination of results of Fremlin-Miller [10], Bukovský-Reclaw-Repický [7], and Bukovský-Haleš [5] implies that a Tychonoff space X satisfies $\mathbf{U}_{\text{fin}}(\mathcal{O}, \Gamma)$ hereditarily if, and only if, X satisfies $\mathbf{U}_{\text{fin}}(\mathcal{O}, \Gamma)$ and is a σ -space (see also [34]).⁸ A similar result was proved for $\mathbf{S}_1(\Gamma, \Gamma)$ in [5]. Problem 7.9 in [7] asks whether every (nice enough) σ -space X satisfying $\mathbf{S}_1(\Omega, \Gamma)$, satisfies $\mathbf{S}_1(\Omega, \Gamma)$ hereditarily. A negative answer was given by Miller [18].

We have, in light of the previous section, a simple reason for the difference between $\mathbf{S}_1(\Omega, \Gamma)$ on one hand, and $\mathbf{S}_1(\Gamma, \Gamma)$ and $\mathbf{U}_{\text{fin}}(\mathcal{O}, \Gamma)$ on the other hand.

⁸The assumptions on X in the cited references are stronger, but: A Tychonoff hereditarily- $\mathbf{U}_{\text{fin}}(\mathcal{O}, \Gamma)$ space cannot have the unit interval $[0, 1]$ as a continuous image, and thus is zero-dimensional. As it is hereditarily Lindelöf, each open set is a countable union of clopen sets, and this suffices for the arguments in the cited references.

Definition 4.1. Let $\mathcal{A} \in \{\Gamma, \Omega, \text{O}, \dots\}$. \mathcal{A} is *Borel superset covering* for X if, for each subspace $Y \subseteq X$ and each $\mathcal{U} \in \mathcal{A}(Y)$, there are $\mathcal{V} \subseteq \mathcal{U}$ and a Borel $B \subseteq X$, such that $Y \subseteq B$ and $\mathcal{V} \in \mathcal{A}(B)$.

Many classical types of covers are Borel superset covering. In particular, we have the following.

Lemma 4.2. Γ and O are Borel superset covering (for all X).

Proof. If $\mathcal{U} \in \Gamma(X)$, take a countable infinite subset $\mathcal{V} = \{U_n : n \in \mathbb{N}\} \subseteq \mathcal{U}$, and $B = \bigcup_m \bigcap_{n \geq m} U_n$. If $\mathcal{U} \in \text{O}(X)$, take $B = \bigcup \mathcal{U}$. \square

By Miller's mentioned result and the following theorem, Ω need not be Borel superset covering.

Theorem 4.3. Let $\mathcal{A}, \mathcal{B} \in \{\Gamma, \Omega, \text{O}\}$, and $\Pi \in \{\text{S}_1, \text{S}_{\text{fin}}, \text{U}_{\text{fin}}\}$. Assume that X is a σ -space, satisfies $\Pi(\mathcal{A}, \mathcal{B})$, and \mathcal{A} is Borel superset covering for X . Then X satisfies $\Pi(\mathcal{A}, \mathcal{B})$ hereditarily.

Proof. Let $Y \subseteq X$ and assume that $\mathcal{U}_1, \mathcal{U}_2, \dots \in \mathcal{A}(Y)$. For each n , pick intermediate Borel sets $Y \subseteq B_n \subseteq X$ such that $\mathcal{U}_n \in \mathcal{A}(B_n)$, and let $B = \bigcap_n B_n$. Then $\mathcal{U}_1, \mathcal{U}_2, \dots \in \mathcal{A}(B)$. As B is Borel and X is a σ -space, B is F_σ in X . By Corollary 2.4, B satisfies $\Pi(\mathcal{A}, \mathcal{B})$, and we can thus obtain from the covers $\mathcal{U}_1, \mathcal{U}_2, \dots$ a cover $\mathcal{V} \in \mathcal{B}(B)$. Then $\mathcal{V} \in \mathcal{B}(Y)$. \square

As for the other direction, we have, by the discussion at the beginning of this section, the following.

Lemma 4.4. If a property P of Tychonoff spaces implies $\text{U}_{\text{fin}}(\text{O}, \Gamma)$ and X satisfies P hereditarily, then X is a σ -space. \square

During his work with the second author on [34], L. Zdomskyy asked whether every subset of \mathbb{R} satisfying $\text{U}_{\text{fin}}(\text{O}, \Omega)$ hereditarily is a σ -space. We show that the answer is “No”, in a very strong sense. We will see that, in the context of Scheepers Diagram, Lemma 4.4 becomes a *criterion*, that is, satisfying P hereditarily implies being a σ -space if, and only if, P implies $\text{U}_{\text{fin}}(\text{O}, \Gamma)$.

We use, in our proof, the method of “forcing”, following an elegant approach of Brendle [4]. A proof avoiding this method (e.g., using the methods of [29]) would be more lengthy and technically involved.

Theorem 4.5. Assume the Continuum Hypothesis. There is $L \subseteq \mathbb{R}$ such that all finite powers of L satisfy $\text{S}_1(\text{B}_\Omega, \text{B}_\Omega)$ hereditarily, but L is not a σ -space.

Proof. Let M_α , $\alpha < \aleph_1$, be an increasing sequence of countable submodels of $H(\lambda)$ (λ large enough) such that:

- (1) $\mathbb{R} \subseteq \bigcup_{\alpha < \aleph_1} M_\alpha$;
- (2) For each $\alpha < \aleph_1$, $M_{\alpha+1} \models M_\alpha$ is countable; and
- (3) $(M_\beta : \beta \leq \alpha) \in M_{\alpha+1}$.

For each α choose a real c_α Cohen generic over M_α . (As M_α is countable, there is such c_α .) Let $L = \{c_\alpha : \alpha < \aleph_1\}$. It is well known that a set L constructed this way is a *Luzin set*, i.e., L is uncountable, but for each meager $M \subseteq \mathbb{R}$, $L \cap M$ is countable.⁹

Lemma 4.6 (folklore). *No Luzin set L is a σ -space.*

Proof. Assume otherwise, and take a countable dense $D \subseteq L$. As $L \setminus D$ is Borel, it is F_σ . Then $D = \bigcap_n U_n$, an intersection of open sets. Then for each n , $L \setminus U_n$ is meager, and thus countable. It follows that $L \setminus D$ is countable, a contradiction. \square

Recall that according to Definition 1.2, X satisfies P^\uparrow if all finite powers of X satisfy P . By [2], or alternatively Lemma 4.2 and Remark 4.9, $S_1(B, B)$ is hereditary. By [26], $S_1(B_\Omega, B_\Omega)^\uparrow = S_1(B, B)^\uparrow$.

Lemma 4.7. *If P is hereditary, then so is P^\uparrow .*

Proof. If $Y \subseteq X$ and X satisfies P^\uparrow , then for each k , $Y^k \subseteq X^k$ and X^k satisfies P . Thus, Y^k satisfies P for all k . \square

Thus, it suffices to prove that all finite powers of L satisfy $S_1(B, B)$. Consider for example L^2 . We have to show (essentially, [26]) that for each Borel $\Psi : \mathbb{R}^2 \rightarrow \mathbb{N}^\mathbb{N}$, there is $g \in \mathbb{N}^\mathbb{N}$ such that for each $(x, y) \in L^2$, $\Psi(x, y)(n) = g(n)$ for some n .

Let $\Psi : \mathbb{R}^2 \rightarrow \mathbb{N}^\mathbb{N}$ be Borel. Ψ is coded by a real. Let α be such that this code belongs to M_α . $\mathbb{N}^\mathbb{N} \cap M_\alpha$ is countable. Take a canonical partition $\mathbb{N} = \bigcup_n I_n$ with each I_n infinite, so that $(I_n : n \in \mathbb{N}) \in M_\alpha$. Enumerate $\mathbb{N}^\mathbb{N} \cap M_\alpha = \{f_n : n \in \mathbb{N}\}$, and take $g \in \mathbb{N}^\mathbb{N}$ such that for each n , $g \upharpoonright_{I_n} = f_n \upharpoonright_{I_n}$. For $x, y \in L$, $\Psi(x, y) \in M_\alpha[x, y]$, an extension of M_α by finitely many Cohen reals, which is in fact either M_α or an extension of M_α by a single Cohen real. As the Cohen forcing does not add eventually different reals, there is n such that $f_n \upharpoonright_{I_n}$ coincides with $\Psi(x, y) \upharpoonright_{I_n}$ on infinitely many values. But $f_n \upharpoonright_{I_n} = g \upharpoonright_{I_n}$. \square

We conclude the section with the following characterization, which may be viewed as a necessary revision of Problem of 7.9 of Bukovský-Reclaw-Repický [7], so that it has a (provably) positive answer.¹⁰

⁹Indeed, each meager set is contained in a meager F_σ set M , which in turn is coded by a single real r . Let α be such that $r \in M_\alpha$. Then for each $\beta > \alpha$, $c_\beta \notin M$.

¹⁰See the discussion at the beginning of the present section.

Theorem 4.8. *A Tychonoff space X satisfies $S_1(\Omega, \Gamma)$ hereditarily if, and only if:*

- (1) X satisfies $S_1(\Omega, \Gamma)$;
- (2) X is a σ -space; and
- (3) Ω is Borel superset covering for X .

Proof. Theorem 4.3 provides the “if” part. For the “only if” part, by Lemma 4.4 it remains to prove (3). Let $Y \subseteq X$, $\mathcal{U} \in \Omega(Y)$. As Y satisfies $S_1(\Omega, \Gamma)$, there are $U_n \in \mathcal{U}$ such that $\{U_n : n \in \mathbb{N}\} \in \Gamma(Y)$. Let $B = \bigcup_m \bigcap_{n \geq m} U_n$. Then $Y \subseteq B$ and $\{U_n : n \in \mathbb{N}\} \in \Omega(B)$. \square

Remark 4.9 (The case of countable Borel covers). In the Borel case, the properties are hereditary for Borel subsets [26], and thus in Theorem 4.3, there is no need to assume that X is a σ -space. This explains why some of the Borel properties are hereditary [2] whereas others are not [18], and why none of the open properties is hereditary [2].

Two major open problems concerning $S_1(B_\Omega, B_\Omega)$ and $S_{\text{fin}}(B_\Omega, B_\Omega)$ are whether these properties are hereditary, and whether they are—like their open variant—preserved by finite powers [2, 32, 18]. By Lemma 4.7, the problems are related: If $S_1(B_\Omega, B_\Omega)$ is preserved by finite powers, then it is equal to $S_1(B, B)^\dagger$, which is hereditary. And similarly for $S_{\text{fin}}(B_\Omega, B_\Omega)$.

Item (2) may be removed from Theorem 4.8 in the Borel case. On the other hand, $U_{\text{fin}}(B, B_\Gamma)$ is hereditary and (for Lindelöf spaces) implies $U_{\text{fin}}(O, \Gamma)$, and thus implies being a σ -set. Theorem 4.5 completes the picture.

5. TOPOLOGICAL GROUPS WITH STRONG COMBINATORIAL PROPERTIES

Problem 10.7 in [32], which is also implicit in Theorem 20 of [29] and in the discussion around it, asks whether the Continuum Hypothesis implies the existence of an uncountable subgroup G of $\mathbb{Z}^\mathbb{N}$ satisfying $S_1(\Omega, \Gamma)$. We will show that the answer is “Yes”, and indeed the weaker hypothesis $\mathfrak{p} = \mathfrak{b}$ suffices to obtain such groups (see the discussion in Section 3). Indeed, Theorem 3.5 gives an uncountable subset of $\mathbb{Z}^\mathbb{N}$ satisfying $S_1(\Omega, \Gamma)$. We will show that for a wide class of properties P , including $S_1(\Omega, \Gamma)$, if X satisfies P , then so is the group generated by X in any Tychonoff topological group G .

There is some restriction on P : Let $F(X)$ be the free topological group generated by X . That is, the group such that each continuous

function f from X into a topological group H , can be extended uniquely to a continuous homomorphism from $F(X)$ to H .¹¹

Lemma 5.1. *Assume that P is a property of Tychonoff topological spaces, which is hereditary for closed subsets and preserved by homeomorphic images. If for each X satisfying P , $F(X)$ satisfies P , then $P = P^\uparrow$.*

Proof. Assume that X satisfies P . Then $F(X)$ satisfies P . For each k , X^k is homeomorphic to $\{x_1 \cdots x_k : x_1, \dots, x_k \in X\}$, a closed subspace of $F(X)$, and thus satisfies P . \square

Thus, we must consider properties P such that $P = P^\uparrow$. For each Q in the Scheepers Diagram, $P = Q^\uparrow$ is as required. By the discussion in the introduction [15]:

- (1) $S_1(O, O)^\uparrow = S_1(\Omega, \Omega)^\uparrow = S_1(\Omega, \Omega)$;
- (2) $S_{\text{fin}}(O, O)^\uparrow = U_{\text{fin}}(O, \Omega)^\uparrow = S_{\text{fin}}(\Gamma, \Omega)^\uparrow = S_{\text{fin}}(\Omega, \Omega)^\uparrow = S_{\text{fin}}(\Omega, \Omega)$.

Moreover, $S_1(\Omega, \Gamma)^\uparrow = S_1(\Omega, \Gamma)$ [11], and $U_{\text{fin}}(O, \Gamma)^\uparrow$ also has a simple characterization [16]. The properties $S_1(\Gamma, \Gamma)^\uparrow$, $S_1(\Gamma, \Omega)^\uparrow$, and $S_1(\Gamma, O)^\uparrow$ seem, however, to be unexplored. By the results of [15], each of these properties is strictly stronger than its non- \uparrow -ed version.

All P in the Scheepers Diagram have the properties required in the following theorem.

Theorem 5.2. *Assume that P is a property of Tychonoff topological spaces, which is hereditary for closed subsets and preserved by continuous images. If P is linearly σ -additive, then for each X satisfying P^\uparrow and each Tychonoff topological group G containing X , the group $\langle X \rangle \leq G$ generated by X satisfies P^\uparrow .*

Proof. It is not difficult to prove the following.

Lemma 5.3. *Assume that P is a property of topological spaces, which is hereditary for closed subsets and preserved by continuous images, and P is linearly σ -additive. Then P^\uparrow also has these three properties. \square*

Fix distinct $a, b \in X \setminus \{1\}$, 1 being the identity element of G . As $\{1, a\}, \{a, b\}$ are discrete subspaces of G , $X \times \{1, a\}$ is homeomorphic to $X \times \{a, b\}$. As X^2 satisfies P^\uparrow , so does its closed subset $X \times \{a, b\}$, and thus so does $X \times \{1, a\}$. Thus, so does the image Y of $X \times \{1, a\}$ under the continuous map $(x, y) \mapsto xy^{-1}$. $Y = X \cup Xa^{-1}$, and as $a \in X$, $1 \in Y$. As $a \in X$, $\langle X \rangle = \langle Y \rangle$. Thus, we may assume that $1 \in X$.

¹¹A thorough introduction to free topological groups is available in [27].

$(\{1\} \times X) \cup (X \times \{1\})$ is a closed subset of X^2 , which satisfies P^\dagger . Thus, $(\{1\} \times X) \cup (X \times \{1\})$ satisfies P^\dagger , and therefore so does its image under the same map, $Y = X \cup X^{-1}$. As $\langle Y \rangle = \langle X \rangle$, we may assume that $X = X^{-1}$, and $1 \in X$.

Thus,

$$\langle X \rangle = \bigcup_n \{x_1 \cdots x_n : x_1, \dots, x_n \in X\}$$

is an increasing union. For each n , $\{x_1 \cdots x_n : x_1, \dots, x_n \in X\}$ is a continuous image of X^n , which satisfies P^\dagger . Thus, $\langle X \rangle$ satisfies P^\dagger . \square

Theorem 5.2 can also be stated in the language of free topological groups. We obtain results analogous (but incomparable) to ones of Banach, Repovš, and Zdomskyy [1].

Theorem 5.4. *Let P be as in Theorem 5.2. For each topological space X satisfying P^\dagger , the free topological group $F(X)$ satisfies P^\dagger .* \square

It follows that for each P in the Scheepers Diagram, and each X satisfying P^\dagger , $\langle X \rangle$ satisfies P^\dagger . Previous constructions of topological groups satisfying P^\dagger for these properties were much more involved than constructions of topological spaces satisfying P^\dagger .

By Theorem 3.5 and Theorem 5.2, we have the following.

Corollary 5.5. *Assume that $\mathfrak{p} = \mathfrak{b}$. There is a subgroup of \mathbb{R} of cardinality \mathfrak{b} , satisfying $\mathbf{S}_1(\Omega, \Gamma)$.* \square

We conclude with an application of the results of this paper. To put it in context, we draw in Figure 2 the Scheepers Diagram, extended to also contain the Borel properties [26]. In the Borel case, some additional equivalences hold, and thus there are fewer distinct properties. In particular, $\mathbf{U}_{\text{fin}}(\mathbf{B}, \mathbf{B}_\Gamma)$, the Borel counterpart of $\mathbf{U}_{\text{fin}}(\mathbf{O}, \Gamma)$, is equivalent to $\mathbf{S}_1(\mathbf{B}_\Gamma, \mathbf{B}_\Gamma)$ [26].

Theorem 5.6. *Assume the Continuum Hypothesis.*

- (1) *There is a subgroup G of \mathbb{R} of cardinality continuum, such that each finite power of G satisfies $\mathbf{S}_1(\mathbf{B}_\Omega, \mathbf{B}_\Omega)$ hereditarily, but G is not a σ -space, and does not satisfy $\mathbf{U}_{\text{fin}}(\mathbf{O}, \Gamma)$.*
- (2) *There is a subgroup H of \mathbb{R} of cardinality continuum, such that all finite powers of H satisfy $\mathbf{S}_1(\mathbf{B}_\Gamma, \mathbf{B}_\Gamma)$ and $\mathbf{S}_{\text{fin}}(\mathbf{B}_\Omega, \mathbf{B}_\Omega)$ hereditarily, but H does not satisfy $\mathbf{S}_1(\mathbf{O}, \mathbf{O})$.*

Proof. By [2], or alternatively Lemma 4.2 and Remark 4.9, $\mathbf{S}_1(\mathbf{B}, \mathbf{B})$, $\mathbf{S}_1(\mathbf{B}_\Gamma, \mathbf{B}_\Gamma)$, and $\mathbf{S}_{\text{fin}}(\mathbf{B}, \mathbf{B})$, are all hereditary. By [26], $\mathbf{S}_1(\mathbf{B}_\Omega, \mathbf{B}_\Omega)^\dagger = \mathbf{S}_1(\mathbf{B}, \mathbf{B})^\dagger$, and similarly for \mathbf{S}_{fin} . Thus, by Lemma 4.7, it suffices to prove the statements with the words “hereditarily” removed.




FIGURE 2. The extended Scheepers Diagram

(1) In the proof of Theorem 4.5, we have constructed a Luzin set $L \subseteq \mathbb{R}$ satisfying $S_1(B_\Omega, B_\Omega)^\uparrow$. L is not a σ -space (Lemma 4.6). Let $G = \langle L \rangle$. By Theorem 5.2, G satisfies $S_1(B_\Omega, B_\Omega)^\uparrow$.

As $L \subseteq G$ and being a σ -space is hereditary, G is not a σ -space. As $S_1(B, B)$ is hereditary and is not satisfied by perfect subsets of \mathbb{R} ,¹² G does not contain any perfect subset of \mathbb{R} . Assume that G satisfies $U_{\text{fin}}(O, \Gamma)$. Theorem 5.5 of [15] tells that sets satisfying $U_{\text{fin}}(O, \Gamma)$ and not containing perfect sets are perfectly meager. In particular, G is meager, and thus so is L , a contradiction.

(2) By Theorem 23 and Lemma 24 of [29], there is a Sierpiński set¹³ $S \subseteq \mathbb{R}$ satisfying $S_1(B_\Gamma, B_\Gamma)^\uparrow$. (This result can alternatively and more easily be proved like Theorem 4.5, by using *random reals* instead of Cohen reals.) Let $H = \langle S \rangle$. By Theorem 5.2, H satisfies $S_1(B_\Gamma, B_\Gamma)^\uparrow$. As $S_1(B_\Gamma, B_\Gamma)$ implies $S_{\text{fin}}(B, B)$, $U_{\text{fin}}(B, B_\Gamma)^\uparrow$ implies $S_{\text{fin}}(B, B)^\uparrow$, which is $S_{\text{fin}}(B_\Omega, B_\Omega)^\uparrow$.

It remains to prove that H does not satisfy $S_1(O, O)$. Assume otherwise. By Remark 4.9, As H satisfies $S_1(B_\Gamma, B_\Gamma)$, it is a σ -space [26]. By Theorem 4.3, its subset S also satisfies $S_1(O, O)$, and thus has Lebesgue measure zero, a contradiction. \square

Acknowledgment. This work is based on a part of the first named author's M.Sc. thesis at the Weizmann Institute of Science, supervised

¹²Assume that a perfect set satisfies $S_1(B, B)$. Then so does the unit interval $[0, 1]$, its continuous image. But even $S_1(O, O)$ implies Lebesgue measure zero.

¹³ $S \subseteq \mathbb{R}$ is *Sierpiński set* if it is uncountable, but has countable intersection with each Lebesgue measure zero set.

by Gady Kozma and Boaz Tsaban. We thank Gady Kozma for stimulating discussions. We owe special thanks to Lyubomyr Zdomsky for reading the paper and making useful comments.

REFERENCES

- [1] T. Banach, D. Repovš, and L. Zdomsky, *σ -Boundedness of free topological groups*, arXiv e-print 0904.1389
- [2] T. Bartoszyński and B. Tsaban, *Hereditary topological diagonalizations and the Menger-Hurewicz Conjectures*, *Proceedings of the American Mathematical Society* **134** (2006), 605–615.
- [3] A. Blass, *Combinatorial cardinal characteristics of the continuum*, in: **Handbook of Set Theory** (M. Foreman, A. Kanamori, and M. Magidor, eds.), Kluwer Academic Publishers, Dordrecht, to appear. <http://www.math.lsa.umich.edu/~ablass/hbk.pdf>
- [4] J. Brendle, *Generic constructions of small sets of reals*, *Topology and its Applications* **71** (1996), 125–147.
- [5] L. Bukovský and J. Haleš, *On Hurewicz properties*, *Topology and its Applications* **132** (2003), 71–79.
- [6] L. Bukovský, I. Reclaw, and M. Repický, *Spaces not distinguishing pointwise and quasinormal convergence of real functions*, *Topology and its Applications* **41** (1991), 25–41.
- [7] L. Bukovský, I. Reclaw, M. Repický, *Spaces not distinguishing convergences of real-valued functions*, *Topology and its Applications* **112** (2001), 13–40.
- [8] K. Ciesielski, A. Millán, and J. Pawlikowski, *Uncountable γ -sets under axiom $\text{CPA}^{\text{game cube}}$* , *Fundamenta Mathematicae* **176** (2003), 143–155.
- [9] M. Džamonja, M. Hrušák, and J. Moore, *Parametrized \diamond principles*, *Transactions of the American Mathematical Society* **356** (2004), 2281–2306.
- [10] D. Fremlin and A. Miller, *On some properties of Hurewicz, Menger and Rothberger*, *Fundamenta Mathematica* **129** (1988), 17–33.
- [11] F. Galvin and A. Miller, *γ -sets and other singular sets of real numbers*, *Topology and its Applications* **17** (1984), 145–155.
- [12] J. Gerlits and Zs. Nagy, *Some properties of $C(X)$, I*, *Topology and its Applications* **14** (1982), 151–161.
- [13] Gary Gruenhage and Paul Szeptycki, *Fréchet-Urysohn for finite sets*, *Topology and its Applications* **151** (2005) 238–259.
- [14] F. Jordan, *There are no hereditary productive γ -spaces*, *Topology and its Applications* **155** (2008), 1786–1791.
- [15] W. Just, A. Miller, M. Scheepers, and P. Szeptycki, *The combinatorics of open covers II*, *Topology and its Applications* **73** (1996), 241–266.
- [16] Lj. Kočinac and M. Scheepers, *Combinatorics of open covers (VII): Groupability*, *Fundamenta Mathematicae* **179** (2003), 131–155.
- [17] Lj. Kočinac, *Selected results on selection principles*, in: **Proceedings of the 3rd Seminar on Geometry and Topology** (Sh. Rezapour, ed.), July 15–17, Tabriz, Iran, 2004, 71–104.
- [18] A. Miller, *A Nonhereditary Borel-cover γ -set*, *Real Analysis Exchange* **29** (2003/4), 601–606.

- [19] A. Miller, *The γ -Borel conjecture*, Archive for Mathematical Logic **44** (2005), 425–434. <http://arxiv.org/math/math.LO/0312308>
- [20] A. Miller, *Cardinal characteristic for relative γ -sets*, Topology and its Applications **156** (2009), 872–878.
- [21] M. Sakai, *Property C'' and function spaces*, Proceedings of the American Mathematical Society **104** (1988), 917–919.
- [22] M. Scheepers, *Combinatorics of open covers I: Ramsey theory*, Topology and its Applications **69** (1996), 31–62.
- [23] M. Scheepers, *$C_p(X)$ and Arhangel'skii's α_i spaces*, Topology and its Applications **89** (1998), 265–275.
- [24] M. Scheepers, *Sequential convergence in $C_p(X)$ and a covering property*, East-West Journal of Mathematics **1** (1999), 207–214.
- [25] M. Scheepers, *Selection principles and covering properties in topology*, Note di Matematica **22** (2003), 3–41.
- [26] M. Scheepers and B. Tsaban, *The combinatorics of Borel covers*, Topology and its Applications **121** (2002), 357–382.
- [27] M. Tkachenko, *Topological groups for topologists: Part II*, Boletín de la Sociedad Matemática Mexicana. Tercera Serie **6** (2000), 1–41.
- [28] B. Tsaban, *Some new directions in infinite-combinatorial topology*, in: **Set Theory** (J. Bagaria and S. Todorcevic, eds.), Trends in Mathematics, Birkhäuser 2006, 225–255.
- [29] B. Tsaban, *\mathfrak{o} -bounded groups and other topological groups with strong combinatorial properties*, Proceedings of the American Mathematical Society **134** (2006), 881–891.
- [30] B. Tsaban, *Additivity numbers of covering properties*, in: **Selection Principles and Covering Properties in Topology** (L. Kočinac, ed.), Quaderni di Matematica 18, Seconda Università di Napoli, Caserta 2006, 245–282.
- [31] B. Tsaban, *A new selection principle*, Topology Proceedings **31** (2007), 319–329.
- [32] B. Tsaban, *Selection Principles and special sets of reals*, in: **Open Problems in Topology II** (E. Pearl, ed.), Elsevier B.V. 2007, 91–108.
- [33] B. Tsaban and L. Zdomskyy, *Combinatorial images of sets of reals and semi-filter trichotomy*, Journal of Symbolic Logic **73** (2008), 1278–1288.
- [34] B. Tsaban and L. Zdomskyy, *Hereditarily Hurewicz spaces and Arhangel'skii sheaf amalgamations*, submitted for publication.

(Orenshtein) DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL

E-mail address: talo@weizmann.ac.il

URL: <http://www.orenshtein.com>

(Tsaban) DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, RAMAT-GAN 52900, ISRAEL; AND DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL

E-mail address: tsaban@math.biu.ac.il

URL: <http://www.cs.biu.ac.il/~tsaban>